

ON POINT ENERGIES, SEPARATION RADIUS AND MESH NORM
FOR s -EXTREMAL CONFIGURATIONS ON
COMPACT SETS IN R^n

By


S.B. Damelin

and

V. Maymeskul

IMA Preprint Series # 2067

(October 2005)



INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS

UNIVERSITY OF MINNESOTA
400 Lind Hall
207 Church Street S.E.
Minneapolis, Minnesota 55455-0436

Phone: 612/624-6066 Fax: 612/626-7370
URL: <http://www.ima.umn.edu>

On Point Energies, Separation Radius and Mesh Norm for s -Extremal Configurations on Compact Sets in \mathbb{R}^n

S. B. Damelin* and V. Maymeskul †

Abstract

We investigate bounds for point energies, separation radius, and mesh norm of certain arrangements of N points on sets A from a class \mathcal{A}^d of d -dimensional compact sets embedded in $\mathbb{R}^{d'}$, $1 \leq d \leq d'$. We assume that these points interact through a Riesz potential $V = |\cdot|^{-s}$, where $s > 0$ and $|\cdot|$ is the Euclidean distance in $\mathbb{R}^{d'}$. With $\delta_s^*(A, N)$ and $\rho_s^*(A, N)$ denoting, respectively, the separation radius and mesh norm of s -extremal configurations, which are defined to yield minimal discrete Riesz s -energy, we show, in particular, the following.

(A) For the d -dimensional unit sphere $S^d \subset \mathbb{R}^{d+1}$ and $s < d - 1$, $\delta_s^*(S^d, N) \geq cN^{-1/(s+1)}$ and, moreover, $\delta_s^*(S^d, N) \geq cN^{-1/(s+2)}$ if $s \leq d - 2$. The latter result is sharp in the case $s = d - 2$. In addition, point energies for s -extremal configurations are asymptotically equal. This observation relates to numerical experiments on observed scar defects in certain biological systems.

(B) For $A \in \mathcal{A}^d$ and $s > d$, $\delta_s^*(A, N) \geq cN^{-1/d}$ and the mesh ratio $\rho_s^*(A, N)/\delta_s^*(A, N)$ is uniformly bounded for a wide subclass of \mathcal{A}^d . We also conclude that point energies for s -extremal configurations have the same order, as $N \rightarrow \infty$.

AMS Classification: Primary 28A78, 52A40; Secondary 31C05, 31C20, 57N16.

Keywords and phrases: Discrete Riesz energy, Extremal configurations, Point energies, Separation radius, Mesh norm, Sphere, Uniform distribution of points on a sphere, Scar defects, Equilibrium measure, Hausdorff measure.

1 Introduction

The problem of distributing a large number of points over the surface of a smooth manifold is an interesting and widely studied problem with numerous applications in diverse areas. To name just a few: spherical t -designs, discrepancy and combinatorics, Monte-Carlo and Quasi-Monte-Carlo methods, approximation theory, finite fields, complexity theory, frame theory, viral morphology, crystallography, molecular structure and electrostatics. We refer the reader to [1]–[11], [13]–[16], [18]–[25], and the many references cited therein for a detailed account of this fascinating subject. In this paper, we are interested in studying point energies, separation and mesh norm for arrangements of N points on a class of d -dimensional compact sets A embedded in $\mathbb{R}^{d'}$. (Here and throughout the paper, $1 \leq d \leq d'$ are integers.) We assume that these N -arrangements interact through the power law (Riesz) potential $V = |\cdot|^{-s}$, where $s > 0$ and $|\cdot|$ is the Euclidean distance in $\mathbb{R}^{d'}$.

Given a compact set $A \subset \mathbb{R}^{d'}$ and a collection $\omega_N = \{x_1, \dots, x_N\}$ of $N \geq 2$ distinct points on A , the discrete Riesz s -energy associated with ω_N is given by

$$E_s(A, \omega_N) := \sum_{1 \leq i < j \leq N} |x_i - x_j|^{-s}. \quad (1.1)$$

*Supported, in part by grants EP/C000285 and NSF-DMS-0439734. S.B. Damelin thanks the Institute for Mathematics and Applications for their hospitality.

†Department of Mathematical Sciences, Georgia Southern University, Statesboro, GA 30460, USA.

E-mail: damelin@ima.umn.edu, vmaymesk@georgiasouthern.edu

Let $\omega_s^*(A, N) := \{x_1^*, \dots, x_N^*\} \subset A$ be a configuration for which $E_s(A, \omega_N)$ attains its minimal value, that is,

$$\mathcal{E}_s(A, N) := \min_{\omega_N \subset A} E_s(A, \omega_N) = E_s(A, \omega_s^*(A, N)).$$

In accordance with convention, we shall call such minimal configurations *s-extremal configurations*. It is well-known that, in general, *s-extremal configurations* are not always unique. For example, in the case of the unit sphere $S^d := \{x \in \mathbb{R}^{d+1} : |x| = 1\}$, they are invariant under rotations.

In this paper, we investigate bounds for point energies, separation radius, and mesh norm of *s-extremal configurations*, which are defined to yield minimal discrete Riesz *s-energy*. With $\delta_s^*(A, N)$ and $\rho_s^*(A, N)$ denoting, respectively, the separation radius and mesh norm of such configurations, we show, in particular, the following. (A) For the *d*-dimensional unit sphere $S^d \subset \mathbb{R}^{d+1}$ and $s < d - 1$, $\delta_s^*(S^d, N) \geq cN^{-1/(s+1)}$ and, moreover, $\delta_s^*(S^d, N) \geq cN^{-1/(s+2)}$ if $s \leq d - 2$. The latter result is sharp in the case $s = d - 2$. In addition, point energies for *s-extremal configurations* are asymptotically equal. This observation relates to numerical experiments on observed scar defects in certain biological systems.

(B) For $A \in \mathcal{A}^d$ and $s > d$, $\delta_s^*(A, N) \geq cN^{-1/d}$ and the mesh ratio $\rho_s^*(A, N)/\delta_s^*(A, N)$ is uniformly bounded for a wide subclass of \mathcal{A}^d . We also conclude that point energies for *s-extremal configurations* have the same order, as $N \rightarrow \infty$.

Natural questions that arise in studying the discrete Riesz energy are:

- (1) What is the asymptotic behavior of $\mathcal{E}_s(A, N)$, as $N \rightarrow \infty$?
- (2) How are *s-extremal configurations* distributed on *A* for large *N*?

It is well-known that answers to these questions essentially depend on the relation between *s* and the Hausdorff dimension $d_H(A)$ of *A*. We demonstrate this fact with the following two classical examples and refer the reader to [15] for more details.

Example 1 The interval $[-1, 1]$, $d_H([-1, 1]) = 1$: In the limiting cases, i.e., $s = 0$ (logarithmic interactions) and $s = \infty$ (best-packing problem), the *s-extremal configurations* are Fekete points and equally spaced points, respectively. It is well-known that Fekete points are distributed on $[-1, 1]$ according to the arcsine measure, which has the density $\mu'_0(x) := (1/\pi)(1 - x^2)^{-1/2}$, while the equally spaced points, $-1 + 2(k - 1)/(N - 1)$, $k = 1, \dots, N$, have the arclength distribution, as $N \rightarrow \infty$. It is also known that $s = 1$ is the critical value in the sense that *s-extremal configurations* are distributed on $[-1, 1]$ differently for $s < 1$ and $s \geq 1$ (see [17, Appendix] and [20]). Indeed, for $s < 1$, the limiting distribution of *s-extremal configurations* has an arcsine-type density

$$\mu'_s(x) := \frac{\Gamma(1 + s/2)}{\sqrt{\pi} \Gamma((1 + s)/2)} (1 - x^2)^{(s-1)/2} \quad (1.2)$$

and, for $s \geq 1$, the limiting distribution is the arclength distribution.

Concerning the minimal energies, they again behave differently for $s < 1$, $s = 1$, and $s > 1$:

$$\mathcal{E}_s([-1, 1], N) \asymp \begin{cases} (1/2)N^2 e_s, & s < 1, \\ (1/2)N^2 \ln N, & s = 1, \\ (1/2)^s \zeta(s) N^{1+s}, & s > 1, \end{cases}$$

where $e_s := [\sqrt{\pi} \Gamma(1 + s/2)] / [\cos(\pi s/2) \Gamma((1 + s)/2)]$ and $\zeta(s)$ stands for the Riemann zeta function.

This dependence of the distribution of *s-extremal configurations* over $[-1, 1]$ and the asymptotics for minimal discrete *s-energy* on *s* can be easily explained from potential theory point of view. Indeed, for a probability Borel measure ν on $[-1, 1]$, its *s-energy* integral is defined to be

$$I_s([-1, 1], \nu) := \iint_{[-1, 1]^2} |x - y|^{-s} d\nu(x) d\nu(y) \quad (1.3)$$

(which can be finite or infinite). Let, for a set of points $\omega_N = \{x_1, \dots, x_N\}$ on $[-1, 1]$,

$$\nu^{\omega_N} := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$$

denote the normalized counting measure of ω_N (so that $\nu^{\omega_N}([-1, 1]) = 1$). Then the discrete Riesz s -energy, associated with ω_N and defined by (1.1), can be written as

$$E_s([-1, 1], \omega_N) = (1/2)N^2 \iint_{x \neq y} |x - y|^{-s} d\nu^{\omega_N}(x) d\nu^{\omega_N}(y), \quad (1.4)$$

where the integral represents a discrete analog of the s -energy integral (1.3) for the point-mass measure ν^{ω_N} .

If $s < 1$, then it is well-known (cf. [17, Appendix]) that the energy integral (1.3) is minimized uniquely by an arcsine-type measure ν_s^* , whose density $\mu'_s(x)$ with respect to the Lebesgue measure is given by (1.2). On the other hand, the normalized counting measure $\nu_{s,N}^*$ of an s -extreme configuration minimizes the discrete energy integral in (1.4) over all configurations ω_N on $[-1, 1]$. Thus one can reasonably expect that, for N large, $\nu_{s,N}^*$ is “close” to ν_s^* and, therefore, the minimal discrete s -energy $\mathcal{E}_s([-1, 1], N)$ is close to $(1/2)N^2 I_s([-1, 1], \nu_s^*) = (1/2)N^2 e_s$.

If $s \geq 1$, then the energy integral (1.3) diverges for every measure ν . Thus, $\mathcal{E}_s([-1, 1], N)$ must grow faster than N^2 . Concerning the distribution of s -extremal points over $[-1, 1]$, the interactions are now strong enough to force them to stay away from each other as far as possible. Of course, depending on s , “far” neighbors still incorporate some energy in $\mathcal{E}_s([-1, 1], N)$, but the closest neighbors are dominating. So, s -extremal points distribute themselves over $[-1, 1]$ in an equally spaced manner.

Example 2 The unit sphere S^d , $d_H(S^d) = d$: Here we again see three distinct cases: $s < d$, $s = d$, and $s > d$. Although it turns out that, for any s , the limiting distribution of s -extremal configurations is given by the normalized area measure on S^d (cf. [17]; [13], [9]; [14], resp.), which is not a big surprise due to the rotation invariance, the asymptotic behavior of $\mathcal{E}_s(S^d, N)$ is quite different. With

$$\tau_{s,d}(N) := \begin{cases} N^2, & s < d, \\ N^2 \ln N, & s = d, \\ N^{1+s/d}, & s > d, \end{cases}$$

it is known that the limit $\lim_{N \rightarrow \infty} \mathcal{E}_s(S^d, N) / \tau_{s,d}(N)$ exists (see [17], [16], [14]). Moreover, in the first two cases, it has the value $(1/2)\gamma_{s,d}$, where

$$\begin{aligned} \gamma_{d,d} &= \frac{\Gamma((d+1)/2)}{d\sqrt{\pi}\Gamma(d/2)}, \\ \gamma_{s,d} &= \frac{\Gamma((d+1)/2)\Gamma(d-s)}{\Gamma((d-s+1)/2)\Gamma(d-s/2)}, \quad s < d. \end{aligned} \quad (1.5)$$

The dependence of the growth rate of $\mathcal{E}_s(S^d, N)$ on s can be explained using potential theory arguments similar to those in Example 1.

2 Class \mathcal{A}^d

In this section, we introduce a class of compact sets $A \subset \mathbb{R}^{d'}$, for which, in the case $s \geq d$ and under some additional assumptions, the asymptotic behavior of $\mathcal{E}_s(A, N)$, separation results, and the limiting distribution of $\omega_s^*(A, N)$ over A (in terms of weak-star convergence of the normalized counting measures) have been recently obtained (see [14, Theorems 2.1–2.4]). We will also give two important properties of sets in \mathcal{A}^d (estimates (2.3) and (2.4) below), which turn out to be crucial in what follows.

For a set $A \subset \mathbb{R}^{d'}$, let $\mathcal{H}^d(A)$ denote its d -dimensional Hausdorff measure (which reduces to d -dimensional Lebesgue measure if $d' = d$).

Definition 2.1 We say that a set A belongs to the class \mathcal{A}^d if, for some $d' \geq d$, $A \subset \mathbb{R}^{d'}$ and

- (1) $\mathcal{H}^d(A) > 0$ and
- (2) A is a finite union of bi-Lipschitz images of compact sets in \mathbb{R}^d , that is

$$A = \bigcup_{i=1}^m \varphi_i(K_i),$$

where each $K_i \subset \mathbb{R}^d$ is compact and $\varphi_i : K_i \rightarrow \mathbb{R}^{d'}$ is bi-Lipschitz on K_i , $i = 1, \dots, m$.

We recall that a mapping $\varphi : K \rightarrow \mathbb{R}^{d'}$ is called bi-Lipschitz (with a constant L) on a compact set $K \subset \mathbb{R}^d$ if there exists a constant L such that for all $x, y \in K$

$$(1/L)|x - y| \leq |\varphi(x) - \varphi(y)| \leq L|x - y|.$$

Clearly, $\mathcal{H}^d(\varphi(K')) \leq L^d \mathcal{H}^d(K')$ for any measurable set $K' \subseteq K$. In particular, it follows that, for any measurable set $A' \subseteq \varphi(K)$,

$$\mathcal{H}^d(A') \leq C [\text{diam}(A')]^d, \quad (2.1)$$

where the constant C depends on L and d only, and $\text{diam}(\cdot)$ means the diameter of a set. Indeed, since $\text{diam}(\varphi^{-1}(A')) \leq L \text{diam}(A')$, the preimage $\varphi^{-1}(A')$ is contained in a ball $B \subset \mathbb{R}^d$ of radius $L \text{diam}(A')$. Thus,

$$\mathcal{H}^d(A') \leq L^d \mathcal{H}^d[\varphi^{-1}(A')] \leq L^d \mathcal{H}^d(B) \leq \frac{2\pi^{d/2} L^{2d}}{d\Gamma(d/2)} [\text{diam}(A')]^d.$$

We now give some examples of sets from the class \mathcal{A}^d .

- Example 3**
- (i) Compact sets A in \mathbb{R}^d with $\mathcal{H}^d(A) > 0$: with $d' = d$, these sets are bi-Lipschitz images of themselves under the identity map. (One can also consider these sets embedded in $\mathbb{R}^{d'}$ for $d' > d$.) For example, balls $B^d(x, r) := \{y \in \mathbb{R}^d : |y - x| \leq r\}$, d -dimensional cubes and parallelepipeds, d -dimensional Cantor sets having positive \mathcal{H}^d -measure.
 - (ii) d -dimensional spheres in \mathbb{R}^{d+1} (more generally, ellipsoids), since a closed hemisphere is a bi-Lipschitz image of a d -dimensional ball under a stereographic projection.
 - (iii) Quasismooth (chord-arc) curves in $\mathbb{R}^{d'}$. These are Jordan curves $A \subset \mathbb{R}^{d'}$ satisfying the following condition: there exists a constant C such that, for any two points $x, y \in A$, the length of the (shortest) subarc of A with endpoints x and y is bounded by $C|x - y|$. In this case, the bi-Lipschitz mapping $\varphi : [0, \text{length}(A)] \rightarrow \mathbb{R}^{d'}$ is given by a natural parametrization of the curve.
 - (iv) Finite unions of sets from (i)–(iii). For instance, a ball $B^d(x, r)$ with a quasismooth outgoing “tail”. Thus, sets in \mathcal{A}^d can have, locally, different Hausdorff dimensions.

We remark the following useful properties of sets $A \in \mathcal{A}^d$. For $r > 0$ and $x \in A$, let

$$E(x, r) := \{y \in A : |y - x| < r\} \quad (2.2)$$

denote a “cap” on A with center x and radius r . Then

$$\mathcal{H}^d(E(x, r)) \leq C_1 r^d \quad (2.3)$$

with a constant C_1 independent of x and r . Indeed, let L be such that all φ_i 's in Definition 2.1 are bi-Lipschitz with the constant L . We note that $\text{diam}(E(x, r)) \leq 2r$ and, using (2.1), conclude that

$$\mathcal{H}^d(E(x, r)) \leq \sum_{i=1}^m \mathcal{H}^d(E(x, r) \cap K_i) \leq mC(2r)^d = C_1 r^d.$$

In particular, $\mathcal{H}^d(A) \leq C_1[\text{diam}(A)]^d < \infty$ since A is compact. Moreover, (2.3) implies that, for $s > d$,

$$\int_{A \setminus E(x,r)} |y-x|^{-s} d\mathcal{H}^d(y) \leq C_2 r^{d-s}, \quad (2.4)$$

where C_2 does not depend on x and r . This inequality can be verified as follows.

$$\begin{aligned} \int_{A \setminus E(x,r)} |y-x|^{-s} d\mathcal{H}^d(y) &= \int_0^\infty \mathcal{H}^d(\{|y-x|^{-s} > t\} \cap \{A \setminus E(x,r)\}) dt \\ &\leq \int_0^{r^{-s}} \mathcal{H}^d(E(x, t^{-1/s})) dt \leq \int_0^{r^{-s}} C_1 t^{-d/s} dt = C_2 r^{d-s}. \end{aligned}$$

3 Main results

Our results concern point energies, separation radius, and the mesh norm for sets $A \in \mathcal{A}^d$ in the case $s > d$. We also obtain new separation estimates for the unit sphere $S^d \subset \mathbb{R}^{d+1}$ in the case $s < d-1$ and show that point energies are asymptotically equal, as $N \rightarrow \infty$.

Throughout the paper, we denote by C, C_1, \dots positive constants, and by c, c_1, \dots sufficiently small positive constants (different each time, in general), that may depend on d, s, A , and other parameters not essential for arguments, but independent of N and other variable quantities.

We define the *point energies* associated with $\omega_s^*(A, N)$ by

$$\mathcal{E}_{j,s}(A, N) := \sum_{\substack{i=1 \\ i \neq j}}^N |x_j^* - x_i^*|^{-s}, \quad j = 1, \dots, N. \quad (3.1)$$

For $s > d$, it was shown in [14] that, if $A \in \mathcal{A}^d$ and $m = 1$ in Definition 2.1 (i.e., $A \subset \mathbb{R}^d$ is a bi-Lipschitz image of one compact set in \mathbb{R}^d), then

$$\mathcal{E}_{j,s}(A, N) \leq CN^{s/d}, \quad 1 \leq j \leq N.$$

We extend this result with

Theorem 3.2 *Let $A \in \mathcal{A}^d$ and $s > d$. Then, for all $1 \leq j \leq N$,*

$$\mathcal{E}_{j,s}(A, N) \leq CN^{s/d}. \quad (3.2)$$

Remark 3.3 Our proof of Theorem 3.2 shows that given d, s , and A in advance, the constant C in (3.2) can be explicitly estimated. Since $\mathcal{E}_s(A, N) = (1/2) \sum_{j=1}^N \mathcal{E}_{j,s}(A, N)$, we conclude that

$$\limsup_{n \rightarrow \infty} \frac{\mathcal{E}_s(A, N)}{N^{1+s/d}} \leq C(d, s, A)$$

with an explicit value of the constant $C(d, s, A)$, and so we obtain an estimate for a constant in the upper bound for the minimal energies.

For $j = 1, \dots, N$ and a set $\omega_N = \{x_1, \dots, x_N\}$ of distinct points on $A \in \mathcal{A}^d$, we let

$$\delta_j(\omega_N) := \min_{i \neq j} \{|x_i - x_j|\}$$

and define

$$\delta(\omega_N) := \min_{1 \leq j \leq N} \delta_j(\omega_N).$$

The quantity $\delta(\omega_N)$ is called the *separation radius* and gives the minimal distance between points in ω_N .

We also define the *mesh norm* $\rho(A, \omega_N)$ of ω_N by

$$\rho(A, \omega_N) := \max_{y \in A} \min_{x \in \omega_N} |y - x|. \quad (3.3)$$

Geometrically, $\rho(A, \omega_N)$ means the maximal radius of a cap $E(y, r)$ (see (2.2)) on A , which does not contain points from ω_N .

These two quantities, $\delta(\omega_N)$ and $\rho(A, \omega_N)$, give a good enough description of the distribution of ω_N over the set A . It is worth mentioning that, even for a sequence $\{\omega_N\}$ of *asymptotically s -extremal* configurations, i.e., configurations satisfying

$$\lim_{N \rightarrow \infty} \frac{E_s(A, \omega_N)}{\mathcal{E}_s(A, N)} = 1,$$

using results from [14], one can get only trivial estimates for the separation radius. Namely,

$$\delta(\omega_N) \geq cN^{-(1/d+1/s)}, \quad s > d.$$

However, for s -extremal configurations on $A \in \mathcal{A}^d$ much better (best possible) estimate for the separation radius holds. As an immediate consequence of Theorem 3.2, we get

Corollary 3.4 *For $A \in \mathcal{A}^d$, $s > d$, and any s -extremal configuration $\omega_s^*(A, N)$ on A ,*

$$\delta_s^*(A, N) := \delta(\omega_s^*(A, N)) \geq cN^{-1/d}. \quad (3.4)$$

Indeed, for any $1 \leq j \leq N$, (3.2) yields

$$cN^{s/d} \geq \mathcal{E}_{j,s}(A, N) \geq [\delta_j(\omega_s^*(A, N))]^{-s},$$

and so $\delta_j(\omega_s^*(A, N)) \geq cN^{-1/d}$ for all j .

Remark 3.5 We remark that (3.4) was proved in [16, Corollary 1.4] for the unit sphere S^d . The same estimate was also shown in [14, Theorem 2.4] for $A \in \mathcal{A}^d$ with an additional condition that $m = 1$ in Definition 2.1. Unfortunately, this latter condition is not satisfied for many sets, including the sphere S^d , and so Corollary 3.4 constitutes an extension of [14, Theorem 2.4].

Separation results for $s < d$ are far more difficult to find in the literature. It trivially follows from Example 2 that

$$\delta(\omega_N) \geq cN^{-2/s}$$

for any sequence $\{\omega_N\}$ of asymptotically s -extremal configurations on the unit sphere S^d . But, even for s -extremal configurations on S^d , there are only two non-trivial results known to the authors: for $d = 2$, $s = 0$ ([10], [22]) and $d \geq 2$, $s = d - 1$ ([8]) it was shown that

$$\delta_s^*(S^d, N) \geq cN^{-1/d}. \quad (3.5)$$

(In [8], two-sided estimates were obtained for more general surfaces.) The estimate (3.5) is quite natural and, intuitively, should be valid for any $s > 0$. A reason for such a lack of results for weak interactions ($s < d$) is that this case require more delicate considerations based on the minimizing property of $\omega_s^*(A, N)$ while strong interactions ($s > d$) prevent points to be very close to each other without affecting the total energy, and separation estimates can be obtained by looking at nearest neighbors only.

Our next result provides a separation estimate in the case $s < d - 1$ for the unit sphere S^d .

Theorem 3.6 For $d \geq 2$ and $s < d - 1$,

$$\delta_s^*(S^d, N) \geq cN^{-1/(s+1)}. \quad (3.6)$$

As a by-product of the proof of Theorem 3.6, we obtain the following

Corollary 3.7 For any $0 < s < d - 1$,

$$\lim_{N \rightarrow \infty} \frac{\max_{1 \leq j \leq N} \mathcal{E}_{j,s}(S^d, N)}{\min_{1 \leq j \leq N} \mathcal{E}_{j,s}(S^d, N)} = 1. \quad (3.7)$$

Indeed, it follows from (4.3), (4.6), and (4.11) that, for N large enough,

$$1 \leq \frac{\max_{1 \leq j \leq N} \mathcal{E}_{j,s}(S^d, N)}{\min_{1 \leq j \leq N} \mathcal{E}_{j,s}(S^d, N)} = \frac{\max_{1 \leq j \leq N} U_j(x_j^*)}{\min_{1 \leq j \leq N} U_j(x_j^*)} \leq 1 + CN^{-1/(s+1)}.$$

Remark 3.8 Numerical computations for a sphere and a torus (see [3], [5], [6], [15]) suggest that, for any $s > 0$, the point energies are nearly equal for almost all points (which are of so called “hexagonal” type). However, some points (“pentagonal”) have elevated energies and some (“heptagonal”) have low energies. The transition from points that are “hexagonal” to those that are “pentagonal” and “heptagonal” induces scar defects, which are conjectured to vanish for N large enough. Thus, Corollary 3.7 confirms this conjecture for $0 < s < d - 1$.

The estimate (3.6) can be improved for $d \geq 3$ and $s \leq d - 2$.

Theorem 3.9 Let $d \geq 3$ and $s \leq d - 2$. Then

$$\delta_s^*(S^d, N) \geq cN^{-1/(s+2)}. \quad (3.8)$$

Note that, when $s = d - 2$, (3.8) gives the best possible estimate (3.5).

Concerning the mesh norm $\rho(A, \omega_N)$, for any sequence $\{\omega_N\}$ of asymptotically s -extremal configurations on A , clearly,

$$\lim_{N \rightarrow \infty} \rho(A, \omega_N) = 0.$$

However, no estimate on the rate of convergence can be made. We show that, for $s > d$, under an additional condition on $A \in \mathcal{A}^d$, the mesh norm and the separation radius of any sequence $\{\omega_s^*(A, N)\}$ of s -extremal configurations on A have the same order, as $N \rightarrow \infty$.

Theorem 3.10 Let $s > d$, $A \in \mathcal{A}^d$, and suppose further that

$$\mathcal{H}^d(E(x, r)) \geq cr^d, \quad (3.9)$$

where a constant $c > 0$ is independent of $x \in A$ and $r > 0$ small enough. Then, for any s -extremal configuration $\omega_s^*(A, N)$ on A ,

$$\rho_s^*(A, N) := \rho(A, \omega_s^*(A, N)) \leq CN^{-1/d}. \quad (3.10)$$

We remark that the condition (3.9) implies the existence of a constant $C_1 > 0$, which depends only on C in (2.1) and c in (3.9), such that

$$\max_{1 \leq j \leq N} \delta_j(\omega_s^*(A, N)) \leq C_1 \rho_s^*(A, N), \quad (3.11)$$

Indeed, let k be an index for which the maximum in (3.11) is attained. Thus,

$$\max_{1 \leq j \leq N} \delta_j(\omega_s^*(A, N)) = \delta_k(\omega_s^*(A, N)) =: \delta^*.$$

The cap $E(x_k, (1/3)\delta^*)$ contains a point y satisfying

$$|y - x_k| \geq \frac{1}{8} \left(\frac{c}{C} \right)^{1/d} \delta^*, \quad (3.12)$$

where constants $C \geq 1$ and $c \leq 1$ are such that (2.1) and (3.9) hold. (Assuming that no such point existed, we would easily conclude that $\text{diam}(E(x_k, (1/3)\delta^*)) \leq (c/C)^{1/d} \delta^*/4$ and so, by (2.1),

$$\mathcal{H}^d(E(x_k, (1/3)\delta^*)) \leq c \left(\frac{\delta^*}{4} \right)^d,$$

which contradicts (3.9).) Since $|y - x_i^*| \geq (2/3)\delta^*$ for all $i \neq k$, taking into account (3.12), we obtain

$$\min_{1 \leq i \leq N} |y - x_i^*| \geq \frac{1}{8} \left(\frac{c}{C} \right)^{1/d} \delta^*,$$

and (3.11) follows.

Corollary 3.4, Theorem 3.10, and (3.11) yield

Corollary 3.11 *Let $s > d$, and assume that $A \in \mathcal{A}^d$ satisfies (3.9). Then, for any s -extremal configuration $\omega_s^*(A, N)$ on A ,*

$$c \leq N^{1/d} \delta_j(\omega_s^*(A, N)) \leq C, \quad 1 \leq j \leq N. \quad (3.13)$$

Since the upper estimate in (3.13) gives

$$\mathcal{E}_{j,s}(A, N) \geq [\delta_j(\omega_s^*(A, N))]^{-s} \geq [CN^{-1/d}]^{-s} = cN^{s/d}, \quad 1 \leq j \leq N,$$

combining this inequality with (3.2), we get

Corollary 3.12 *For $s > d$, $A \in \mathcal{A}^d$ satisfying (3.9), and any s -extremal configuration $\omega_s^*(A, N)$ on A , there holds*

$$1 \leq \frac{\max_{1 \leq j \leq N} \mathcal{E}_{j,s}(A, N)}{\min_{1 \leq j \leq N} \mathcal{E}_{j,s}(A, N)} \leq C. \quad (3.14)$$

Remark 3.13 Corollary 3.12 says that, for $s > d$ and a set $A \in \mathcal{A}^d$ satisfying (3.9), the point energies are asymptotically of the same order, as $N \rightarrow \infty$. Most likely, this is the best possible assertion in the sense that an analog of (3.7) does not hold, in general.

Remark 3.14 Simple examples show that the upper estimates in (3.13) and (3.14) are not valid, in general, for a set $A \in \mathcal{A}^d$ without an additional condition on its geometry. For instance, if $x \in \mathbb{R}^{d+1}$ and $|x| > 1$, then $A := S^d \cup \{x\}$, clearly, does not enjoy these properties.

The measure condition (3.9) in Theorem 3.10 can be omitted if $m = 1$ in Definition 2.1 (in particular, if $A \subset \mathbb{R}^d$ is a compact set with positive d -dimensional Lebesgue measure).

Theorem 3.15 *Let $A \in \mathcal{A}^d$ and suppose that A is the image of one compact set $K \subset \mathbb{R}^d$ under a bi-Lipschitz mapping φ . Then (3.10) holds.*

The remainder of this paper is devoted to proofs of our results.

4 Proofs

First, we introduce some notations and properties of s -extreme configurations, which will be used in subsequent proofs.

Let $A \in \mathcal{A}^d$, $\omega_s^*(A, N) = \{x_1^*, \dots, x_N^*\}$ be an s -extremal configuration on A ,

$$\nu_{s,N}^* := \frac{1}{N} \sum_{i=1}^N \delta_{x_i^*}$$

denote the normalized counting measure of $\omega_s^*(A, N)$, and let

$$U^{\nu_{s,N}^*}(x) := \int_{\mathbb{R}^{d'}} |x-y|^{-s} d\nu_{s,N}^*(y) = \int_A |x-y|^{-s} d\nu_{s,N}^*(y) = \frac{1}{N} \sum_{i=1}^N |x-x_i^*|^{-s} \quad (4.1)$$

be the potential associated with $\nu_{s,N}^*$. We also introduce functions

$$U_j(x) := U^{\nu_{s,N}^*}(x) - \frac{1}{N} |x-x_j^*|^{-s} = \frac{1}{N} \sum_{\substack{i=1 \\ i \neq j}}^N |x-x_i^*|^{-s} = \int_A |x-y|^{-s} d\nu_{s,j,N}^*(y), \quad (4.2)$$

where $\nu_{s,j,N}^* := \nu_{s,N}^* - (1/N)\delta_{x_j^*}$. Note that the total mass of $\nu_{s,j,N}^*$ is $(N-1)/N$. It follows from (3.1) that

$$\mathcal{E}_{j,s}(A, N) = NU_j(x_j^*). \quad (4.3)$$

Since $\omega_s^*(A, N)$ is an s -extremal configuration, $U_j(x)$ attains its minimal value at x_j^* . Thus, for any $x \in A$ and $1 \leq j \leq N$, we have

$$U^{\nu_{s,N}^*}(x) = U_j(x) + \frac{1}{N} |x-x_j^*|^{-s} \geq U_j(x_j^*) + \frac{1}{N} |x-x_j^*|^{-s} = \frac{1}{N} [\mathcal{E}_{j,s}(A, N) + |x-x_j^*|^{-s}]. \quad (4.4)$$

Therefore,

$$\sum_{j=1}^N U^{\nu_{s,N}^*}(x) \geq \frac{1}{N} \sum_{j=1}^N \mathcal{E}_{j,s}(A, N) + \frac{1}{N} \sum_{j=1}^N |x-x_j^*|^{-s} = \frac{2}{N} \mathcal{E}_s(A, N) + U^{\nu_{s,N}^*}(x)$$

and so

$$U^{\nu_{s,N}^*}(x) \geq \frac{2}{N(N-1)} \mathcal{E}_s(A, N) > \frac{2}{N^2} \mathcal{E}_s(A, N), \quad x \in A. \quad (4.5)$$

4.1 Proof of Theorem 3.2

Our proof will follow from (2.3), (2.4), and ideas used in [16, Section 5]. For the reader's convenience, we present a sketch of the proof.

Let $A \in \mathcal{A}^d$, $\omega_s^*(A, N) = \{x_1^*, \dots, x_N^*\}$ be an s -extremal configuration on A , and let $U_j(x)$ be defined by (4.2), $j = 1, \dots, N$. We denote $D_i := A \setminus E(x_i^*, [2NC_1/\mathcal{H}^d(A)]^{-1/d})$, where C_1 is the constant from (2.3), and $D := \cap_{i=1}^N D_i$. Then, using (2.4), we get

$$\begin{aligned} \int_D U_j(x) d\mathcal{H}^d(x) &= \frac{1}{N} \sum_{\substack{i=1 \\ i \neq j}}^N \int_D |x-x_i^*|^{-s} d\mathcal{H}^d(x) \leq \frac{1}{N} \sum_{\substack{i=1 \\ i \neq j}}^N \int_{D_i} |x-x_i^*|^{-s} d\mathcal{H}^d(x) \\ &\leq \frac{1}{N} \sum_{\substack{i=1 \\ i \neq j}}^N C_2 \left[\frac{2NC_1}{\mathcal{H}^d(A)} \right]^{(s-d)/d} < C_2 \left[\frac{2C_1}{\mathcal{H}^d(A)} \right]^{(s-d)/d} N^{s/d-1}. \end{aligned}$$

On the other hand, by (2.3),

$$\mathcal{H}^d(D) = \mathcal{H}^d(A) - \mathcal{H}^d\left(\bigcup_{i=1}^N E\left(x_i^*, [2NC_1/\mathcal{H}^d(A)]^{-1/d}\right)\right) \geq \mathcal{H}^d(A) - \sum_{i=1}^N C_1[2NC_1/\mathcal{H}^d(A)]^{-1} = \frac{\mathcal{H}^d(A)}{2},$$

and so the minimizing property of $\omega_s^*(A, N)$ yields

$$U_j(x_j^*) \leq \frac{1}{\mathcal{H}^d(D)} \int_D U_j(x) d\mathcal{H}^d(x) \leq (C_1 C_2)^{(s-d)/d} [\mathcal{H}^d(A)/2]^{-s/d} N^{s/d-1} = C_3 N^{s/d-1}.$$

(Note that, since C_1 and C_2 depend explicitly on d , s , and A , so does C_3 .)

Finally, using (4.3), we obtain (3.2). ■

4.2 Proof of Theorems 3.6 and 3.9

Note that, since $d' = d + 1$ and $s < d - 1 = d' - 2$, the potential

$$U^\nu(x) := \int_{\mathbb{R}^{d+1}} |x - y|^{-s} d\nu(y)$$

is superharmonic in \mathbb{R}^{d+1} for any finite positive Borel measure ν on \mathbb{R}^{d+1} (see [17, Thm. 1.4]).

To save in writing, we will use a standard notation σ for the normalized d -dimensional Hausdorff measure \mathcal{H}^d on S^d .

Let $\gamma_{s,d}$ be defined by (1.5). It is well-known that (for any $0 < s < d$) $\sigma(x)$ is the equilibrium measure for S^d , and so $U^\sigma(x) \equiv \gamma_{s,d}$ on S^d . Using this fact and integrating both sides of

$$\frac{1}{N} \sum_{i=1}^N |x - x_i^*|^{-s} = U^{\nu_{s,N}}(x) \geq \frac{1}{N} [\mathcal{E}_{j,s}(A, N) + |x - x_j^*|^{-s}] = U_j(x_j^*) + \frac{1}{N} |x - x_j^*|^{-s}$$

(see (4.1), (4.4), and (4.3)) against $\sigma(x)$, we get

$$\gamma_{s,d} \geq U_j(x_j^*) + \frac{1}{N} \gamma_{s,d}.$$

Thus,

$$U_j(x_j^*) \leq \frac{N-1}{N} \gamma_{s,d} < \gamma_{s,d}, \quad j = 1, \dots, N. \quad (4.6)$$

Next, we obtain estimates for $U_j(x_j^*)$ from below. It was shown in ([16], [25]) that

$$\frac{1}{2} \gamma_{s,d} N^2 - C_1 N^{1+\alpha(s,d)} \leq \mathcal{E}_s(S^d, N) \leq \frac{1}{2} \gamma_{s,d} N^2 - c_1 N^{1+s/d}, \quad (4.7)$$

$$\alpha(s, d) = \begin{cases} s/(s+2), & 0 < s < d-2, \\ s/d, & d-2 \leq s < d \end{cases}$$

For $d-2 \leq s < d$, (4.7) gives the exact bounds for the second term in the asymptotic behavior of $\mathcal{E}_s(S^d, N)$. In the case $0 < s < d-2$, the lower estimate in (4.7), most likely, is not best possible.

We now use estimates (4.5) and (4.7) to conclude that, for $x \in S^d$,

$$U^{\nu_{s,N}}(x) > \frac{2}{N^2} \left(\frac{1}{2} \gamma_{s,d} N^2 - C_1 N^{1+\alpha(s,d)} \right) = \gamma_{s,d} - C_2 N^{\alpha(s,d)-1}. \quad (4.8)$$

Since $U^{\nu_{s,N}^*}(x)$ is superharmonic, by the minimal value principle (4.8) also holds for $|x| \leq 1$.

Next, fixed $\varepsilon_0 > 0$ small enough, let $0 < \varepsilon < \varepsilon_0$. For any x satisfying $1 < |x| \leq 1 + \varepsilon$, we denote $\tilde{x} := (2 - |x|x)/|x|$. It can be easily verified that

$$\min_{y \in S^d} \frac{|\tilde{x} - y|}{|x - y|} = \frac{3 - |x|}{|x| + 1} = 1 - \frac{2(|x| - 1)}{|x| + 1} > 1 - \varepsilon$$

(with the minimum attained at $y = -x/|x|$). Therefore, using (4.8), we get

$$\begin{aligned} U^{\nu_{s,N}^*}(x) &= N^{-1} \sum_{i=1}^N |x - x_i^*|^{-s} = N^{-1} \sum_{i=1}^N |\tilde{x} - x_i^*|^{-s} \left(\frac{|\tilde{x} - x_i^*|}{|x - x_i^*|} \right)^s \\ &\geq (1 - \varepsilon)^s U^{\nu_{s,N}^*}(\tilde{x}) \geq (1 - C_s \varepsilon) \left(\gamma_{s,d} - C_2 N^{\alpha(s,d)-1} \right) \geq \gamma_{s,d} - C_2 N^{\alpha(s,d)-1} - C_3 \varepsilon. \end{aligned}$$

Thus,

$$U^{\nu_{s,N}^*}(x) \geq \gamma_{s,d} - C_2 N^{\alpha(s,d)-1} - C_3 \varepsilon, \quad |x| \leq 1 + \varepsilon. \quad (4.9)$$

We now apply this estimate to establish Theorems 3.6 and 3.9. Let us assume that, for some $\delta > 0$, i and j , $|x_i^* - x_j^*| \leq (1/2)\delta$, and consider the function

$$U_{i,j}(x) := U_j(x) - N^{-1} |x - x_i^*|^{-s} = U^{\nu_{s,N}^*}(x) - N^{-1} \left(|x - x_j^*|^{-s} + |x - x_i^*|^{-s} \right) = N^{-1} \sum_{k \neq i,j} |x - x_k^*|^{-s}.$$

Then it follows from (4.9) that, for any x satisfying $|x - x_j^*| = \delta$ and $|x| \leq 1 + \varepsilon$,

$$\begin{aligned} U_{i,j}(x) &\geq \gamma_{s,d} - C_2 N^{\alpha(s,d)-1} - C_3 \varepsilon - N^{-1} \left[\delta^{-s} + \left(\frac{\delta}{2} \right)^{-s} \right] \\ &= \gamma_{s,d} - C_2 N^{\alpha(s,d)-1} - C_3 \varepsilon - C_4 N^{-1} \delta^{-s} \end{aligned} \quad (4.10)$$

with constants C_2 , C_3 , and C_4 independent of ε , δ , N , and x .

We begin with the proof of (3.6). Choosing $\varepsilon = \delta = N^{-1/(s+1)}$, (4.10) yields for $|x - x_j^*| = N^{-1/(s+1)}$

$$U_{i,j}(x) \geq \gamma_{s,d} - C_2 N^{\alpha(s,d)-1} - C_3 N^{-1/(s+1)} - C_4 N^{-1} N^{s/(s+1)} \geq \gamma_{s,d} - C_5 N^{-1/(s+1)}.$$

(Here, we have used the fact that, for $s < d - 1$, $1 - \alpha(s,d) > 1/(s+1)$.) Clearly, $U_{i,j}(x)$ is superharmonic in \mathbb{R}^{d+1} , and so we conclude that

$$U_{i,j}(x_j^*) \geq \gamma_{s,d} - C_5 N^{-1/(s+1)}.$$

Therefore

$$U_j(x_j^*) = U_{i,j}(x_j^*) + N^{-1} |x_i^* - x_j^*|^{-s} \geq \gamma_{s,d} - C_5 N^{-1/(s+1)} + N^{-1} |x_i^* - x_j^*|^{-s} \quad (4.11)$$

Combining (4.6) and (4.11), we finally get

$$\gamma_{s,d} > \gamma_{s,d} - C_5 N^{-1/(s+1)} + N^{-1} |x_i^* - x_j^*|^{-s}.$$

Hence,

$$C_5 N |x_i^* - x_j^*|^s \geq N^{1/(s+1)},$$

and (3.6) is proved. \blacksquare

For (3.8), let $P(x_j^*)$, denote the d -dimensional hyperplane touching S^d at x_j^* . Then, if $x \in P(x_j^*)$ with $|x - x_j^*| = \delta > 0$ (i.e., x belongs to the $(d-1)$ -dimensional sphere with center x_j^* and radius δ contained in

$P(x_j^*)$), then clearly $|x| < 1 + \delta^2$. Thus, choosing $\delta = N^{-1/(s+2)}$ and $\varepsilon = \delta^2 = N^{-2/(s+2)}$, we conclude from (4.10) that

$$U_{i,j}(x) \geq \gamma_{s,d} - C_2 N^{s/(s+2)-1} - C_3 N^{-2/(s+2)} - C_4 N^{-1} N^{s/(s+2)} = \gamma_{s,d} - C_6 N^{-2/(s+2)}$$

on the $(d-1)$ -dimensional sphere $\{|x - x_j^*| = N^{-1/(s+2)}\} \cap P(x_j^*)$. Since $s \leq d-2$, it is easy to see that $U_{i,j}(x)$, restricted to $P(x_j^*)$, is superharmonic and, therefore,

$$U_{i,j}(x_j^*) \geq \gamma_{s,d} - C_6 N^{-2/(s+2)}.$$

Similarly to (4.11), we find that

$$U_j(x_j^*) = U_{i,j}(x_j^*) + N^{-1} |x_i^* - x_j^*|^{-s} \geq \gamma_{s,d} - C_6 N^{-2/(s+2)} + N^{-1} |x_i^* - x_j^*|^{-s},$$

which together with (4.6) yields

$$\gamma_{s,d} > \gamma_{s,d} - C_6 N^{-2/(s+2)} + N^{-1} |x_i^* - x_j^*|^{-s}.$$

Thus,

$$C_6 N |x_i^* - x_j^*|^s \geq N^{2/(s+2)},$$

and (3.8) follows. ■

4.3 Proof of Theorems 3.10 and 3.15

To simplify notations, let us denote $\rho^* := \rho_s^*(A, N)$. We can obviously assume that

$$\rho^* \geq cN^{-1/d}, \tag{4.12}$$

where c is the constant from the separation estimate (3.4).

In [14, Lemma 3.1], the authors proved a two-sided estimate on $\mathcal{E}_s(A, N)$ for the case when $A \subset \mathbb{R}^d$ is a bounded set with nonempty interior. (See also [23, Theorem 2] for the case of the unit sphere $S^d \subset \mathbb{R}^{d+1}$.) It can be easily verified that the approach used in proving the lower estimate,

$$\mathcal{E}_s(A, N) \geq c_1 N^{1+s/d}$$

with a constant $c_1 > 0$ depending on d, s , and A , but not on N , can be applied to any set $A \in \mathcal{A}^d$. Therefore, for the potential $U^{\nu_{s,N}^*}(x)$ defined in (4.1), (4.5) yields the estimate

$$U^{\nu_{s,N}^*}(x) > \frac{2}{N^2} \left(c_1 N^{1+s/d} \right) = 2c_1 N^{s/d-1}, \quad x \in A.$$

In particular, this estimate is valid for $x = y^*$, where y^* is a point at which the maximum in (3.3) is attained (with $\omega_N = \omega_s^*(A, N)$). Thus

$$U^{\nu_{s,N}^*}(y^*) > 2c_1 N^{s/d-1}. \tag{4.13}$$

We now derive an upper estimate for $U^{\nu_{s,N}^*}(y^*)$. First, we note that

$$|y^* - x_i^*| \geq \rho^*, \quad i = 1, \dots, N.$$

Let $E_i^* := E(x_i^*, (c/4)N^{-1/d})$. It is clear that

$$E_i^* \cap E_j^* = \emptyset, \quad i \neq j. \tag{4.14}$$

We also have that

$$\rho^* = \min_{x \in \omega_s^*(A, N)} |x - y^*| \leq |x_i^* - y^*|, \quad 1 \leq i \leq N. \tag{4.15}$$

Then for any $x \in E_i^*$, $1 \leq i \leq N$, (4.12) and (4.15) yield

$$|x - y^*| \leq |x - x_i^*| + |x_i^* - y^*| \leq \frac{cN^{-1/d}}{4} + |x_i^* - y^*| \leq \frac{\rho^*}{4} + |x_i^* - y^*| \leq \frac{5}{4} |x_i^* - y^*|,$$

which implies that

$$|x_i^* - y^*|^{-s} \leq \left(\frac{5}{4}\right)^s \min_{x \in E_i^*} \left\{ |x - y^*|^{-s} \right\}, \quad (4.16)$$

and

$$|x - y^*| \geq |x_i^* - y^*| - |x - x_i^*| \geq |x_i^* - y^*| - \frac{cN^{-1/d}}{4} \geq |x_i^* - y^*| - \frac{\rho^*}{4} \geq \frac{3\rho^*}{4}. \quad (4.17)$$

Now using (4.16), (3.9), and (3.4), we get

$$\begin{aligned} |x_i^* - y^*|^{-s} &\leq \left(\frac{5}{4}\right)^s \min_{x \in E_i^*} \left\{ |x - y^*|^{-s} \right\} \leq \left(\frac{5}{4}\right)^s \frac{1}{\mathcal{H}^d(E_i^*)} \int_{E_i^*} |x - y^*|^{-s} d\mathcal{H}^d(x) \\ &\leq \frac{C_1}{[(c/4)N^{-1/d}]^d} \int_{E_i^*} |x - y^*|^{-s} d\mathcal{H}^d(x) \leq C_2 N \int_{E_i^*} |x - y^*|^{-s} d\mathcal{H}^d(x). \end{aligned} \quad (4.18)$$

By (4.17),

$$\bigcup_{i=1}^N E_i^* \subset A \setminus E(y^*, 3\rho^*/4) =: D.$$

Thus (4.18), (4.14), and (2.4) yield

$$\begin{aligned} U^{\nu_{s,N}^*}(y^*) &= N^{-1} \sum_{i=1}^N |x_i^* - y^*|^{-s} \leq C_2 \sum_{i=1}^N \int_{E_i^*} |x - y^*|^{-s} d\mathcal{H}^d(x) \\ &\leq C_2 \int_D |x - y^*|^{-s} d\mathcal{H}^d(x) \leq C_3 \left(\frac{3\rho^*}{4}\right)^{d-s} = C_4 (\rho^*)^{d-s}. \end{aligned} \quad (4.19)$$

Combining (4.13) and (4.19), we obtain $2c_1 N^{s/d-1} \leq C_4 (\rho^*)^{d-s}$, and (3.10) follows. \blacksquare

Theorem 3.15 can be proved in a very similar way. The only difference is that, this time, we can obtain the upper estimate (4.19) without the condition (3.9) by using the mapping φ . Indeed, let $z_i^* := \varphi^{-1}(x_i^*)$, $t^* := \varphi^{-1}(y^*)$. Since φ is bi-Lipschitz (say, with a constant L), we have

$$|x_i^* - y^*| \geq (1/L) |z_i^* - t^*|.$$

Therefore,

$$U^{\nu_{s,N}^*}(y^*) \leq N^{-1} \sum_{i=1}^N [(1/L) |z_i^* - t^*|]^{-s} = L^s N^{-1} \sum_{i=1}^N |z_i^* - t^*|^{-s}. \quad (4.20)$$

We also note that

$$\min_{j \neq i} |z_j^* - z_i^*| \geq \frac{c}{L} N^{-1/d}, \quad (4.21)$$

where c is the constant from (3.4), and (see (4.15))

$$|z_i^* - t^*| \geq \frac{\rho^*}{L}, \quad 1 \leq i \leq N. \quad (4.22)$$

Let B_i^* denote the ball in \mathbb{R}^d with center z_i^* and radius $(cN^{-1/d})/(4L)$. It follows from (4.21) that

$$B_i^* \cap B_j^* = \emptyset, \quad i \neq j. \quad (4.23)$$

Then, for $z \in B_i^*$, $1 \leq i \leq N$, (4.22) yields

$$|z - t^*| \leq \frac{cN^{-1/d}}{4L} + |z_i^* - t^*| \leq \frac{\rho^*}{4L} + |z_i^* - t^*| \leq \frac{5}{4} |z_i^* - t^*|,$$

which implies that

$$|z_i^* - t^*|^{-s} \leq \left(\frac{5}{4}\right)^s \min_{z \in B_i^*} \left\{ |z - t^*|^{-s} \right\}, \quad (4.24)$$

and

$$|z - t^*| \geq |z_i^* - t^*| - \frac{cN^{-1/d}}{4L} \geq |z_i^* - t^*| - \frac{\rho^*}{4L} \geq \frac{3\rho^*}{4L}. \quad (4.25)$$

Using (4.24), we conclude that

$$|z_i^* - t^*|^{-s} \leq \left(\frac{5}{4}\right)^s \frac{1}{\mathcal{H}^d(B_i^*)} \int_{B_i^*} |z - t^*|^{-s} d\mathcal{H}^d(z) \leq C_1 N \int_{B_i^*} |z - t^*|^{-s} d\mathcal{H}^d(z). \quad (4.26)$$

By (4.25),

$$\bigcup_{i=1}^N B_i^* \subset \mathbb{R}^d \setminus B^d(t^*, 3\rho^*/(4L)) =: D.$$

Thus (4.20), (4.26), and (4.23) yield

$$\begin{aligned} U^{\nu_s^*, N}(y^*) &\leq L^s N^{-1} \sum_{i=1}^N |z_i^* - t^*|^{-s} \leq C_1 L^s \sum_{i=1}^N \int_{B_i^*} |z - t^*|^{-s} d\mathcal{H}^d(z) \\ &\leq C_1 L^s \int_D |z - t^*|^{-s} d\mathcal{H}^d(z) = C_2 \left(\frac{3\rho^*}{4L}\right)^{d-s} = C_3 (\rho^*)^{d-s}. \end{aligned}$$

Combining this estimate with (4.13), we obtain (3.10). ■

References

- [1] M. Atiya and P. Sutcliffe, *The geometry of point particles*, R. Soc. Lond. Proc., Ser. A, Math. Phys. Eng. Sci., **458** (2002): 1089–1115.
- [2] B. Bajnok, S. B. Damelin, J. Li, and G. Mullen, *A finite field algorithm for distributing points on a d-dimensional sphere*, Computing (Springer), **68** (2002): 97–109.
- [3] A. R. Bausch, M. J. Bowick, A. Cacciuto, A. D. Dinsmore, M. F. Hsu, M. G. Nikolaides, A. Travesset, and D. A. Weitz, *Grain Boundary Scars and Spherical Crystallography*, Science, March 14 (2003): 1716–1718.
- [4] J. Benedetto and M. Fickus, *Finite normalized tight frames*, Adv. Comput. Math., **18** (2003): 357–385.
- [5] M. Bowick, A. Cacciuto, D. R. Nelson, and A. Travesset, *Crystalline order on a sphere and the generalized Thomson problem*, Phys. Rev. Lett., **89** (2002): 185502.

- [6] M. Bowick, D. R. Nelson, and A. Travesset, *Curvature-Induced defect unbinding in Toroidal Geometries*, To appear in Phys. Rev. E, (2003). Available at: arXiv.org/cond-mat/0309709
- [7] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der Mathematischen Wissenschaften, **290**, Springer Verlag, New York, 1999.
- [8] B. E. J. Dahlberg, *On the distribution of Fekete points*, Duke Math. J., **45** (1978): 537–542.
- [9] S. B. Damelin and P. Grabner, *Energy functionals, numerical integration and asymptotic equidistribution on the sphere*, Numerical integration and its complexity (Oberwolfach, 2001), J. Complexity **19** (2003): 231–246; *Corrigendum to “Energy Functionals, Numerical Integration and Asymptotic Equidistribution on the Sphere*, J. of Complexity, **20** (2004): 883–884.
- [10] P. Dragnev, D. A. Legg and D. W. Townsend, *On the separation of logarithmic points on the sphere*, Approximation Theory X: Abstract and Classical Analysis, Charles K. Chui, Larry L. Schumaker, and Joachim Stoeckler (eds.), Vanderbilt University Press 2002: 137–144.
- [11] T. Erber and G. M. Hockney, *Equilibrium configurations of N equal charges on the sphere*, Adv. Chem. Phys., **98** (1991): L1369–L1377.
- [12] M. Fekete, *Über Interpolation*, Z. Angew. Math. Mech., **6** (1926): 410–413.
- [13] M. Götz and E. B. Saff, *Note on d -extremal configurations for the sphere in \mathbb{R}^{d+1}* , In: Recent progress in multivariate approximation, Internat. Ser. Numer. Math., Birkhauser, Basel, **137** (2001): 159–162.
- [14] D. Hardin and E. B. Saff, *Minimal Riesz energy point configurations for rectifiable d -dimensional manifolds*, Advances in Math, to appear.
- [15] D. Hardin and E. B. Saff, *Discretizing manifolds via minimum energy points*, Notices of the AMS, November 2004, pp. 1186–1194.
- [16] A. B. J. Kuijlaars and E. B. Saff, *Asymptotics of minimal discrete energy on the sphere*, Trans. Amer. Math. Soc., **350** (1998): 523–538.
- [17] N. S. Landkof, *Foundations of Modern Potential Theory*, Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, Berlin, 1972.
- [18] A. Lubotzky, R. Phillips, and P. Sarnak, *Hecke operators and distributing points on the sphere I*, Comm. Pure App. Math., **39** (1986): S148–S186.
- [19] A. Lubotsky, R. Phillips, and P. Sarnak, *Hecke operators and distributing points on the sphere II*, Comm. Pure App. Math. **39** (1986): S401–S420.
- [20] A. Martinez-Finkelshtein, V. V. Maymeskul, E. A. Rakhmanov, and E. B. Saff, *Asymptotics for minimal discrete Riesz energy on curves in \mathbb{R}^d* , Canad. J. Math. **56** (2004): 529–552.
- [21] E. A. Rakhmanov, E. B. Saff, and Y. M. Zhou, *Minimal discrete energy on the sphere*, Mathematical Research Letters, **1** (1994): 647–662.
- [22] E. A. Rakhmanov, E. B. Saff, and Y. M. Zhou, *Electrons on the sphere*, In: Computational Methods and Function Theory (R. M. Ali, St. Ruscheweyh, and E. B. Saff, Eds.), World Scientific Publ., 1995: 111–127.
- [23] E. B. Saff and A. B. J. Kuijlaars, *Distributing many points on a sphere*, Mathematical Intelligencer, **19** (1997): 5–11.
- [24] S. Smale, *Mathematical problems for the next century*, Mathematical Intelligencer, **20** (1998): 7–15.
- [25] G. Wagner, *On the means of distances on the surface of a sphere (lower bounds)*, Pacific J. Math., **144** (1990): 389–398.